

International Foundation for Functional Gastrointestinal Disorders

IFFGD 700 W. Virginia St., #201 Milwaukee, WI 53204

Phone: 414-964-1799 Toll-Free (In the U.S.): 888-964-2001

Free (In the U.S.): 888-964-2001 Fax: 414-964-7176

Internet: www.iffgd.org

(239)

© Copyright 2009-2014 by the International Foundation for Functional Gastrointestinal Disorders

Is Constipation and Bloating Related to Menstrual Periods?

By: Peter Whorwell, M.D., Ph.D., Professor of Medicine and Gastroenterology, Wythenshawe Hospital, Education and Research Centre, Manchester, United Kingdom

QUESTION – For the last 5 months or so, I have experienced severe constipation and bloating pain a couple of days after my period; it would last 4–5 days and go away. I tried Metamucil and it does not help. I use Midol or Aleve for pain during my menstrual period, and I try to keep the dosage to no more than 2–3 pills during the 5 days. Any suggestions?

ANSWER – Many healthy women experience some change of bowel function at the time of their periods. The problem usually lasts only a few days. If severe, an antispasmodic may help for cramping. If you become very constipated, a laxative is perfectly acceptable.

Not surprisingly, women with irritable bowel syndrome (IBS) suffer even more trouble at this time of the month. One abnormality that quite frequently occurs in patients with IBS is that the bowel is over-sensitive and as a result reacts more to things that upset it. For instance, something that might cause a tummy upset for a couple of days in a person without IBS could cause trouble for a couple of weeks in a person with IBS.

If you have a diagnosis of IBS, you need to be a little careful with anti-inflammatory medicines such as Aleve (or others such as Advil, Motrin, etc.) as this type of drug can occasionally upset IBS. Drugs related to acetaminophen (such as Tylenol, Endocet, etc.) are much better for routine pain relief in IBS patients. Be sure to follow the directions on the drug label. Always

let your doctor know what medicines you take, whether prescription or over-the-counter.

The symptoms around the time of your period that you describe can sometimes be related to another gastrointestinal disorder or a gynecological problem. It is important to always tell your doctor when you experience new symptoms or a noticeable change in chronic symptoms.

Suggested Reading

Heitkemper, M.M. *Gynecological Aspects of Irritable Bowel Syndrome*. IFFGD. Fact Sheet No. 123.

Schmulson, M. *Understanding Bloating and Distension*. IFFGD Fact Sheet No. 262.

The information in this fact sheet was adapted from the Clinical corner section of Digestive Health Matters, Vol. 16, No. 1.

Opinions expressed are an author's own and not necessarily those of the International Foundation for Functional Gastrointestinal Disorders (IFFGD). IFFGD does not guarantee or endorse any product in this publication nor any claim made by an author and disclaims all liability relating thereto.

This article is in no way intended to replace the knowledge or diagnosis of your doctor. We advise seeing a physician whenever a health problem arises requiring an expert's care.

IFFGD is a nonprofit education and research organization. Our mission is to inform, assist, and support people affected by gastrointestinal disorders. For more information, or permission to reprint this article, write to IFFGD, 700 W. Virginia St., #201, Milwaukee, WI 53204. Toll-free (in the U.S.): 888-964-2001 or 414-964-1799. Visit our websites at: www.aboutibs.org or www.aboutconstipation.org.